



BCB 2,4-D LV6

Broadleaf Herbicide

FOR SELECTIVE CONTROL OF MANY BROADLEAF WEEDS IN CEREAL GRAINS (WHEAT, BARLEY, MILLET, OATS, AND RYE), CORN, [RED] POTATOES, SORGHUM, SOYBEAN (PREPLANT), FALLOW CROPLAND, FORESTS, GRASS PASTURES, RANGELAND, CONSERVATION RESERVE PROGRAM ACRES, ORNAMENTAL TURF GRASS (INCLUDING TURFGRASS GROWN FOR SOD OR SEED), NON-CROPLAND. ALSO, FOR CONTROL OF TREES BY INJECTION.

ACTIVE INGREDIENT:	% By Weight
2,4-D, 2-ethylhexyl ester*	87.6%
OTHER INGREDIENTS	12.4%
TOTAL	100.0%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid.....58.1%, 5.5 lbs./gal.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

EPA Reg. No.: 104199-3

EPA Est. No.: 82614-IND-001

Manufactured for:



Better Choice Brands LLC
4252 E Log Providence Road
Columbia, MO 65201

NET CONTENTS: MB 265

FIRST AID	
IF IN EYES	• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED	• Call a poison control center or doctor immediately for treatment. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR CLOTHING	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center at 1-800-222-1222 or doctor, or going for treatment. In case of an emergency involving this product, call CHEMTREC at 1-800-424-9300 .	

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION / PRECAUCIÓN**

Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical-resistant gloves made of barrier laminate, nitrile rubber ≥ 14 mils, butyl rubber ≥ 14 mils, or Viton ≥ 14 mils, and
- Chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements.

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements: Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides 40 CFR 170.607(f). When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides 40 CFR 170.607(d-f), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users Should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Product should not be used in or near greenhouses.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, protective eyewear, chemical-resistant gloves made of barrier laminate, nitrile rubber $\geq$ 14 mils, butyl rubber $\geq$ 14 mils, or Viton $\geq$ 14 mils, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

PRODUCT INFORMATION

This product contains 2-ethylhexyl ester of 2,4-D. In cropland, this herbicide is for controlling hard-to-kill weeds, bindweed, thistle, smartweeds, wild garlic, curled dock, tansy ragwort and wild onions.

INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT.

Crop varieties vary in response to 2,4-D and some are easily injured. Apply this herbicide to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties or local use situations that may affect crop tolerance to 2,4-D, consult your seed company, state Agricultural Extension Service or qualified crop consultant for advice.

WEED RESISTANCE

Any weed population may contain plants that are naturally resistant to 2,4-D, the active ingredient in this product, and to other herbicides with the same mode of action. ATTENTION: These resistant weed biotypes will not be controlled by this product. Consult advisors such as your local agricultural extension service for agronomic management practices to minimize the occurrence of 2,4-D resistance and considerations for supplemental control measures.

Weed Management

To minimize the occurrence of 2,4-D resistant biotypes, observe the following general weed management practices:

- Scout application site before and after herbicide applications.
- Start with a clean application site, using either a burndown herbicide application or tillage.
- Control weeds early when they are relatively small.
- Add other herbicides (e.g. a selective and/or a residual herbicide) and cultural practices (e.g. tillage or crop rotation) where appropriate.
- Utilize the specified label rate for the most difficult to control weed in your field. Avoid tank mixtures with other herbicides that reduce this product's efficacy (through antagonism), or tank mixture directions that encourage application rates of this product below the label directions.
- Control weed escapes and prevent weeds from setting seeds.
- Clean equipment before moving from field to field to minimize the spread of weed seed or plant parts.
- Report any incidence of repeated non-performance of this product on a particular weed to your BETTER CHOICE BRANDS LLC representative, local retailer, or county extension agent.

Management of 2,4-D-Resistant Biotypes

Since the occurrence of new 2,4-D weeds cannot be determined until after product use and scientific confirmation, manufacturer is not responsible for any losses that may result from the failure of this product to control 2,4-D resistant weed biotypes.

The following good agronomic practices are recommended to reduce the spread of confirmed 2,4-D resistant biotypes:

- If a naturally occurring resistant biotype is present in your application site, this product should be tank-mixed or applied sequentially with an appropriately labeled herbicide with a different mode of action to achieve control.
- Cultural and mechanical control practices (e.g. crop rotation or tillage) may also be used as appropriate.
- Scout treated application site after herbicide applications and control escaping weeds including resistant biotypes before they set seed.
- Thoroughly clean equipment before leaving fields known to contain resistant biotypes.

PRODUCT RESTRICTIONS

- Do not apply this product through any type of irrigation system.
- Do not use in or near a greenhouse.
- Do not use the same spray equipment for other purposes unless thoroughly cleaned. Crops contacted by sprays or spray drift may be killed or suffer significant stand loss with extensive quality and yield reduction.
- Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

MIXING INSTRUCTIONS

Add 1/2 the required amount of water to the spray tank, then add this product with agitation, and finally, the balance of the water with continued agitation. This material forms an emulsion in water, not a solution. This tends to separate on standing. Provide agitation to prevent such separation and insure uniform spray mixtures.

COMPATIBILITY

If this herbicide is to be tank mixed with fertilizers or with other pesticides, test compatibility prior to mixing. To test for compatibility, use a small container and mix a small amount (0.5 to 1 quart) of spray, combining all ingredients in the same ratio as the anticipated use. If any indications of physical incompatibility develop, do not use this mixture for spraying. Indications of incompatibility usually will appear within 5 to 15 minutes after mixing.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

COMBINATION WITH LIQUID NITROGEN FERTILIZER

Use 0.33 to 0.5 pint of this product per acre for weeding and feeding corn, small grains or grass pastures as directed on this label. Use fertilizer at rates recommended by supplier or Extension Service Specialist.

Fill the spray tank about half full with the liquid fertilizer, then add this product with vigorous agitation, and complete filling the tank with fertilizer. Apply immediately and continue agitation in the spray tank during application. Application during very cold weather (near freezing) is not advisable. Do not allow mixture to stand overnight. Incompatibility may be encountered with some fertilizer brands or under some environmental conditions. If in doubt, test a small sample in the dilution ratio planned for application.

NOTE: Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentration will reduce the hazard of leaf burn.

APPLICATION INSTRUCTIONS

Apply with calibrated air or ground equipment using sufficient spray volume to provide adequate coverage of target weeds or as otherwise directed in specific use directions.

For aerial application - Do not apply less than 2 gallon total spray volume per acre.

For ground applications - Do not apply less than 10 gallon total spray volume per acre.

The higher spray volumes will be needed under the following circumstances:

- For difficult to control vegetation,
- For large vegetation,
- Under conditions where control is more difficult, or
- When tank mixing with oils.

Rate Ranges and Application Timing

The lower dosages given will be satisfactory for young, succulent growth of sensitive weed species. For less sensitive species and under conditions where control is more difficult, the higher dosages will be needed. Apply this product during warm weather when weeds are young and actively growing.

Spot Treatments

To prevent misapplication, apply spot treatments with a calibrated boom or with hand sprayers using a fixed spray volume per 1,000 square feet as indicated below.

Hand-Held Sprayers: Hand-held sprayers may be used for spot applications of this product. Apply the spray uniformly and at a rate equivalent to a broadcast application. Application rates in the table are based upon the application rate for an area of 1,000 square feet. Mix the amount of this product (fl oz or mL) corresponding to the desired broadcast rate in 1 to 3 gallons of spray. To calculate the amount of product required for larger areas, multiply the table value (fl oz or mL) by the thousands of square feet to be treated.

Rate Conversion Table for Spot Treatment:

Label Broadcast Rate (pint/acre)							
1/2	2/3	3/4	1	2	3	4	5
Equivalent Amount of BCB 2,4-D LV6 per 1000 sq ft							
1/5 fl oz ¹ (5.9 mL)	1/4 fl oz (7.4 mL)	1/3 fl oz (9.9 mL)	3/8 fl oz (11 mL)	3/4 fl oz (22 mL)	1 fl oz (30 mL)	1-1/2 fl oz (44 mL)	3 fl oz (89 mL)

¹ Conversion factors: 1 fl oz = 29.6 (30) mL

Band Application

BCB 2,4-D LV6 may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

Band width in inches	X	Broadcast rate	=	Band rate
Row width in inches		per acre		acre treated acre
Band width in inches	X	Broadcast volume	=	Band volume
Row width in inches		per acre		acre treated acre

Volumetric % Product in Spray Solution Table

Pints	0.50	0.67	0.75	1.00	2.00	3.00	4.00	5.00
Gallons	0.06	0.08	0.09	0.13	0.25	0.38	0.50	0.63
2.00	3.13%	4.17%	4.69%	6.25%	12.50%	18.75%	25.00%	31.25%
3.00	2.08%	2.78%	3.13%	4.17%	8.33%	12.50%	16.67%	20.83%
4.00	1.56%	2.08%	2.34%	3.13%	6.25%	9.38%	12.50%	15.63%
5.00	1.25%	1.67%	1.88%	2.50%	5.00%	7.50%	10.00%	12.50%
6.00	1.04%	1.39%	1.56%	2.08%	4.17%	6.25%	8.33%	10.42%
7.00	0.89%	1.19%	1.34%	1.79%	3.57%	5.36%	7.14%	8.93%
8.00	0.78%	1.04%	1.17%	1.56%	3.13%	4.69%	6.25%	7.81%
9.00	0.69%	0.93%	1.04%	1.39%	2.78%	4.17%	5.56%	6.94%
10.00	0.63%	0.83%	0.94%	1.25%	2.50%	3.75%	5.00%	6.25%
12.50	0.50%	0.67%	0.75%	1.00%	2.00%	3.00%	4.00%	5.00%
15.00	0.42%	0.56%	0.63%	0.83%	1.67%	2.50%	3.33%	4.17%
20.00	0.31%	0.42%	0.47%	0.63%	1.25%	1.88%	2.50%	3.13%
25.00	0.25%	0.33%	0.38%	0.50%	1.00%	1.50%	2.00%	2.50%
30.00	0.21%	0.28%	0.31%	0.42%	0.83%	1.25%	1.67%	2.08%
50.00	0.13%	0.17%	0.19%	0.25%	0.50%	0.75%	1.00%	1.25%
100.00	0.06%	0.08%	0.09%	0.13%	0.25%	0.38%	0.50%	0.63%

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application (e.g., ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a Coarse or coarser spray, apply only as a Coarse or coarser spray (ASABE S572).

When applying sprays that contain 2,4-D mixed with other active ingredients that require a medium or more fine spray, apply only as a Medium or coarser spray (ASABE S572).

Wind Speed

Do not apply at wind speeds greater than 15 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, non-target plants within residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this by adjusting the path of the aircraft upwind.

Do not apply with a nozzle height greater than 4 feet above the crop canopy.

2,4-D esters may volatilize during conditions of low humidity and high temperatures. Do not apply during conditions of low humidity and high temperatures.

WEEDS LIST

ANNUAL AND BIENNIAL WEEDS

Annual yellow sweet clover	Flixweed	Poorjoe	Vetches
Beggarticks ¹	Geranium, Carolina ^[3]	Prickly lettuce ^{1,[3]}	Wild carrot ^{1,[3]}
Bittercress, smallflowered ^[3]	Hemp, wild	Purslane, common ^[3]	Wild lettuce ^[3]
Bitterweed	Hairy galinsoga	Pusley, Florida	Wild parsnip
Bull thistle	Jewelweed	Ragweed (common, giant)	Wild radish
Buttercup, smallflowered ^{1,[3]}	Jimsonweed	Rape, wild	
Carpetweed	Knotweed ^{1,[3]}	Rocket, yellow	
Cinquefoil, common ^[3]	Kochia	Rough fleabane	
Cinquefoil, rough ^[3]	Lambsquarters, common	Russian thistle ¹	
Coffeeweed	Lupines	Salsify (western or common)	

Common broomweed*
Common burdock
Common cocklebur
Copperleaf, Virginia^[3]
Croton (Texas or woolly)
Evening primrose (common or cutleaf)

Mallow (Venice, dwarf, little)
Marestail^[3]
Marshelder
Morningglory (common, ivy, woolly)
Mousetail^[3]
Mustard (except blue mustard)
Pennycress (fanweed)
Pepperweed (except perennial)^[3]
Pigweeds² (*Amaranthus* spp.)

Shepherdspurse
Sicklepod
Smartweeds^{1,[3]} (annual species)
Sneezeweed, bitter
Sowthistle (annual or spiny)
Spanishneedles
Sunflower
Tansy mustard
Thistle, bull
Thistle, musk¹
Thistle, Russian (tumbleweed)¹
Velvetleaf
Vervains^{1,[3]}

PERENNIAL WEEDS

Alfalfa^{1,[3]}
Austrian fieldcress¹
Bindweed^{1,[3]} (hedge, field & European)
Blue lettuce
Bluweed, Texas
Broomweed
Bullnettle^{1,[3]}
Canadian thistle^{1,[3]}
Carrot, wild¹
Catnip
Chicory^[3]
Clover, red^{1,[3]}
Coffeeweed
Cutleaf^[3]
Dandelion

Docks¹
Dogbanes¹
Evening primrose
Goldenrod¹
Ground ivy¹
Hawkweed¹ (orange)
Healall
Hoary cress¹

Jerusalem – artichoke
Loco, bigbend
Many-flowered aster¹
Nettles (including stinging)
Pennywort
Plantains
Ragwort, tansy¹
Sowthistle (perennial)
Tansy ragwort¹
Vervains¹

Western ironweed^{1,[3]}
Wild garlic¹
Wild onion¹
Wormwood

¹ These species may require repeat applications and/or use of the higher rates on this product label even under ideal conditions for application.

² Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

^[3] This product may not be used to control this weed in the state of California.]

SPECIFIC USE DIRECTIONS

CEREAL GRAINS (WHEAT, BARLEY, MILLET, OATS, and RYE)

(Not underseeded with legumes)

APPLICATION TIMING	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Post-emergence Annual and biennial weeds Perennial broadleaf weeds	0.33 to 1.33 pints* 0.66 to 1.33 pints*	Apply after grain is well tillered (usually about 4 to 8 inches high) but before boot. Do not spray grain in boot to dough stage.
Wild onion or garlic	1 to 1.33 pints*	Apply 1 pint when grain is well tillered and wild garlic or onion plants are small. Apply 1.33 pints after harvest in the crop stubble. For control of new fall growth of wild onion or garlic, refer to FALLOWLAND use directions.

Emergency Weed Control in Wheat Perennial broadleaf weeds	1.75 pints*	Apply when weeds are approaching bud stage, but do not spray grain during the boot to dough stage . The 1.75 pints per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury.
Preharvest	0.66 pint*	Apply when grains are in the hard dough stage to suppress large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to induce succulent weed growth.
*Do not apply to grain in the seedling stage. Maximum rate 1 pint per acre from jointing to before boot stage. Treatment at this growth stage (jointing to before boot) should be used where such increased risk of injury to crop is acceptable. Higher rates listed on this label are more likely to cause crop injury.		

PRECAUTIONS FOR USE ON CEREAL GRAINS

The higher rates (greater than 1.0 pints per acre) increase the risk of grain injury and should be used only where the weed control problem justifies the risk of grain damage.

RESTRICTIONS FOR USE ON CEREAL GRAINS

- The preharvest interval (PHI) is 14 days.
- Do not apply to grain in the seedling stage.
- Postemergence
 - Limited to one postemergence application per crop cycle.
 - Maximum of 1.75 pints (1.25 lbs 2,4-D ae) per acre per application.
- Preharvest
 - Limited to one preharvest application per crop cycle.
 - Maximum of 0.66 pint (0.5 lb 2,4-D ae) per acre per application.
- Limited to 2.5 pints (1.75 lb 2,4-D ae) per acre per crop cycle.
- For aerial application, apply this product in 2 or more gallons of water per acre.
- For ground application, use a minimum of 10 gallons of water per acre.
- Do not mix with oil for crop uses.
- Do not permit dairy animals or meat animals being finished for slaughter to forage treated grain fields within 2 weeks after treatment.
- Do not feed treated straw to livestock if a preharvest treatment as described above is applied.

TANK MIXTURES FOR CEREAL GRAINS

BCB 2,4-D LV6 and Products Containing Bromoxynil* on Wheat, Barley, Millet, Oats, and Rye

GEOGRAPHIC LOCATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Cereal All Areas Except: Washington, Oregon, Idaho, Colorado, Wyoming and Montana	0.33 to 0.75 pint Plus a bromoxynil product at specified label rates	Bromoxynil will control some annual weeds that are resistant to this product and may be tank mixed for broader spectrum weed control on cereal grains.

Washington, Oregon, Idaho, Colorado, Wyoming and Montana	0.33 to 0.75 pint Plus a bromoxynil product at specified label rates	TO PREPARE THE SPRAY: First mix this product in water then add the bromoxynil product. Use the higher rates for larger weeds or where weed growth is slow due to dry or cold weather. Apply before weeds are 6 inches high. Use 10 to 25 gallons total spray volume per acre with ground equipment or use 2 to 10 gallons total spray volume with air application, or sufficient spray solution for adequate coverage. Use higher volume on larger weeds.
--	---	--

*It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

CORN (FIELD, POPCORN AND SWEET)

APPLICATION TIMING / STAGE OF GROWTH	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Preplant (Burndown)	0.75 to 1.33 pints	To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn. Apply 7 to 14 days before planting. Use high rate for control of less susceptible weeds or cover crops, weeds in advanced stages of development, or under less favorable growth conditions.
Preemergence	0.75 to 1.33 pints	Apply any time after planting, but before corn emerges to control broadleaf weed seedlings or existing cover crops. Use high rate for control of less susceptible weeds or cover crops, weeds in advanced stages of development, or under less favorable growth conditions.
Postemergence Annual and biennial weeds Crop up to 8 inches tall Crop 8 inches tall to tasseling (direct spray only) Perennial broadleaf weeds	0.33 to 0.66 pint 0.66 pint 0.75 pint	Avoid spraying just after corn leaves unfold, as injury may occur. Apply when weeds are small and corn is less than 8 inches tall (to top of crop canopy). If corn more than 8 inches tall must be sprayed, use drop nozzles and directed spray to keep spray off foliage. Treat perennial weeds are in the bud to bloom stage. Do not spray when corn is in tassel to hard dough stage.
Preharvest (Field Corn and Popcorn Only)	2 pints	After the hard dough (denting) stage, to suppress weeds that interfere with harvest, bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower and velvetleaf, and to decrease production of weed seeds, spray with air or ground equipment. The high rate will be needed for tough weeds under stress. Do not apply preharvest to sweet corn.

PRECAUTIONS FOR USE ON FIELD CORN, POPCORN AND SWEET CORN

- Preplant or preemergence applications to light sandy soils is not recommended.
- Corn hybrids vary in response to 2,4-D and some are easily injured. Contact the seed company or your Agricultural Experiment Station or Extensional Weed Specialist for this information.
- Corn treated with 2,4-D may exhibit stem brittleness for 8 to 10 days following application. During this period, the crop is more susceptible to stem breakage from wind or cultivation.

RESTRICTIONS FOR USE ON FIELD CORN AND POPCORN

- The preharvest interval (PHI) is 7 days.
- Do not use treated crop as fodder for 7 days following application.
- Preplant or Preemergence
 - Limited to one preplant or preemergence application per crop cycle.
 - Maximum of 1.33 pints (1 lb 2,4-D ae) per acre per application.
- Postemergence
 - Limited to one postemergence application per crop cycle.
 - Maximum of 0.66 pint (0.5 lb 2,4-D ae) per acre per application.
- Preharvest
 - Limited to one postharvest application per crop cycle.
 - Maximum of 2.18 pints (1.5 lb 2,4-D ae) per acre per application.
- Limited to 4.36 pints (3 lb 2,4-D ae) per acre per crop cycle.

RESTRICTIONS FOR USE ON SWEET CORN

- The preharvest interval (PHI) is 45 days.
- Do not use treated crop as fodder for 7 days following application.
- Minimum of 21 days between applications.
- Preplant or Preemergence
 - Limited to one preplant or preemergence application per crop cycle.
 - Maximum of 1.33 pints (1 lb 2,4-D ae) per acre per application.
- Postemergence
 - Limited to one postemergence application per crop cycle.
 - Maximum of 0.75 pint (0.5 lb 2,4-D ae) per acre per application.
- Limited to 2.18 pints (1.5 lb ae) per acre per crop cycle.

FALLOWLAND AND CROP STUBBLE

Fallowland is idle land, postharvest to crops or between crops.

TYPE OF WEEDS	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
FALLOWLAND AND CROP STUBBLE Annual broadleaf weeds	1 to 1.33 pints	Use the lower rate when weeds are small (2 to 3 inches tall) and growing actively. Use a higher in the rate range when weeds are larger and under less favorable growth conditions.
Biennial broadleaf weeds	1.33 to 2.75 pints	Spray while musk thistles or other biennial species are in the seedling to rosette stage, and before flower stalks are initiated. The lower rate can be used in spring during rosette stage. In fall or after flower stalks have developed, use the highest rate.
Perennial broadleaf weeds	1.33 to 2.9 pints	Spray weeds in bud to bloom stage, or in good vegetative growth. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead.
Wild garlic and onion in crop stubble	2.75 to 2.9 pints	Apply to new regrowth of wild onion or garlic which occurs in the fall following harvest of cereal grains, soybeans, corn or grain sorghum.

PRECAUTION FOR USE ON FALLOWLAND

- For best weed control results, do not cultivate for at least 2 weeks after application or until top growth is dead.

RESTRICTIONS AND LIMITATIONS FOR USE ON FALLOWLAND

- The preharvest interval (PHI) is 7 days.
- Minimum of 30 days between applications.

- Limited to 5.8 pints (4.0 lb 2,4-D ae) per acre per use season.
- Maximum of 2.9 pints (2.0 lb 2,4-D ae) per acre per application.
- Do not apply more than two times per year.

Planting in Treated Areas

Labeled Crops: Within 29 days after an application of this product, plant only those crops listed on this or other registered 2,4-D labels. Follow more stringent limitations, if any, provided in directions for specific crops. Labeled crops may be at risk of crop injury or loss if planted soon after application, especially during the first 14 days. Degradation factors described below should be considered in weighing this risk.

Other Crops: All other crops may be planted 30 days or more after application without concern for illegal residues in the planted crop. However, under certain conditions, there may be a risk of injury to susceptible crops. Degradation factors described below should be considered in weighing this risk. Under normal conditions, any crop may be planted without risk of injury if at least 90 days of soil temperatures above freezing have elapsed since application.

Degradation Factors: When planting into treated areas, the risk of crop injury is less if lower rates of product were applied and conditions following application have included warm, moist soil conditions that favor rapid breakdown of 2,4-D. Risk is greater if higher rates of product were applied and soil temperatures have been cold and/or soils have been excessively wet or dry in the days following application. Consult your local agricultural extension service or information about susceptible crops and typical conditions in your area.

[RED] POTATOES (Only for Use on [Red] Potatoes Intended for Fresh Market)

APPLICATION TIMING	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Postemergence	1.65 fl. oz.	[Red] Potatoes: Properly timed applications of this product generally enhance red color, aid in storage retention of red color, improve skin appearance, increase tuber set, and improve tuber size uniformity (fewer jumbos). Crop response may vary depending on variety, stress factors, and local conditions. Varieties with naturally dark red color generally benefit less from treatment.

PRECAUTIONS FOR USE ON [RED] POTATOES

- Make first application when potatoes are in the pre-bud stage (about 7 to 10 inches high) and make a second application about 10 to 14 days later.
- Consult with Agricultural Extension Service and other qualified crop advisors for local recommendations.
- The specific spray volume selected should be sufficient for good coverage of plants.

RESTRICTIONS FOR USE ON [RED] POTATOES

- The preharvest interval (PHI) is 45 days.
- Minimum of 10 days between applications.
- Do not apply more than two times per year.
- Postemergence
 - Limited to two postemergence applications per crop cycle.
 - Maximum of 1.65 fluid ounces (0.07 lb 2,4-D ae) per acre per application.
- Apply 1.65 fluid ounces of this product per acre in 2 to 25 gallons of water using ground or aerial equipment.

SORGHUM [Grain Sorghum (Milo) and Forage Sorghum]

APPLICATION TIMING / STAGE OF GROWTH	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Postemergence Crop 6-8 inches tall	0.33 – 0.75 pint	Apply when sorghum is 6 to 15 inches tall. If sorghum is more than 8 inches tall (to top of crop canopy), use drop nozzles and apply as a directed spray to keep spray off foliage. Do not treat during the boot, flowering or early dough stages.
Crop 8-15 inches tall (directed spray only)	0.5 to 0.75 pint	

PRECAUTIONS FOR USE ON SORGHUM

- Temporary crop injury can be expected under conditions of high soil moisture and high air temperatures. If it is necessary to apply this product under these conditions, use no more than 2/3 pint per acre.
- Sorghum hybrids vary in tolerance to 2,4-D. Some are easily injured. Apply only to varieties known to be tolerant to 2,4-D. Consult the seed company or your agricultural experiment station or extension service weed specialist for this information.

RESTRICTIONS FOR USE ON SORGHUM

- The preharvest interval (PHI) is 30 days.
- Do not permit meat or dairy animals to consume treated crop as fodder or forage for 30 days following application.
- Postemergence
 - Limited to one postemergence application per crop cycle.
 - Maximum of 0.75 pint (0.5 lb 2,4-D ae) per acre per application.
- **Do not treat during the boot, flowering or early dough stages.**

SOYBEAN (PREPLANT ONLY)

APPLICATION TIMING	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Preplant (Burndown)	0.5 to 0.66 pint	Apply not less than 7 days prior to planting soybeans, when weeds are small and actively growing. Use the higher rate on larger weeds and when perennials are present. Some weeds may require repeat treatment for adequate control (see WEED LIST and below). Maximum of 0.66 pint per preplant application. Limited to 2 preplant applications per crop cycle.
	0.66 to 1.33 pints	Apply not less than 15 days prior to planting soybeans, when weeds are actively growing. Limited to 1 application per crop cycle. Apply no more than 1.33 pints of this product in one season prior to planting soybeans. In addition to those weeds found on the WEED LIST, this product will suppress or control the following broadleaf weeds frequently encountered in reduced tillage soybean production systems: bull nettle, smallflowered bittercress, Carolina geranium, smallflowered buttercup, common and rough cinquefoil, red clover*, horseweed or marestalk, mousetail, wild mustard, field pennycress, cutleaf evening primrose, common purslane, speedwell, and Virginia copperleaf. * These weeds are only partially controlled. After applying, plant soybean seed as deep as practical or at least 1-1/2 to 2 inches deep. Adjust the planter press

		wheel, if necessary, to ensure that planted seed is completely covered. If desired, this product may be applied pre-plant to soybeans in tank mixtures with other herbicides* containing the following active ingredients, glyphosate, isopropylamine salt; glyphosate, potassium salt; glufosinate; glufosinate and sodium salt of fomesafen, metribuzin and chlorimuron; tribenuron-methyl and chlorimuron; and flumioxazin; and others that are registered for pre-plant soybean use.
--	--	--

*It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

PRECAUTIONS FOR USE IN SOYBEANS (PREPLANT)

- Unacceptable injury to soybeans planted in fields previously treated with this product may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather and agronomic factors such as the amount of weed vegetation and previous crop residue present that may be in effect between the time of this product's application and the emergence of the soybean plant.

RESTRICTIONS FOR USE IN SOYBEANS (PREPLANT)

- If choosing multiple preplant applications per crop cycle:
 - Limited to 2 preplant applications per crop cycle.
 - Maximum of 0.66 pint (0.5 lb 2,4-D ae) per acre per preplant application.
 - Apply no less than 7 days prior to planting soybeans.

OR

- If choosing a single preplant application per crop cycle:
 - Limited to 1 preplant application per crop cycle.
 - Maximum of 1.5 pints (1.0 lb 2,4-D ae) per acre per preplant application.
 - Apply no less than 15 days prior to planting soybeans.

ADDITIONAL RESTRICTIONS FOR USE IN SOYBEANS (PREPLANT)

- Do not apply this product when weather conditions such as temperature, air inversions, or wind favor drift from treated areas to susceptible plants.
- Do not apply this herbicide prior to planting soybeans if you are not prepared to accept the results of soybean injury including possible loss of stand and yield.
- Do not replant fields treated with this herbicide in the same growing season with crops other than those labeled for 2,4-D preplant use.
- Do not mow or cultivate weeds prior to treating with this product as poor control may result.
- Do not cut for feed treated hay, forage, or fodder or graze treated soybeans to livestock.
- Only one application of this product may be made prior to planting soybeans per growing season.
- Do not cut for feed or graze treated cover crops to livestock.

FORESTRY, RANGELAND, ESTABLISHED PASTURE, AND NON-CROPLAND AREAS

FORESTRY

Forest site preparation, forest roadsides, brush control, established conifer release (including Christmas trees and reforestation areas)

TREATMENT SITE / METHOD OF APPLICATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Annual weeds	1.33 to 2.75 pints	Apply when weeds are small and actively growing, before the bud stage. Apply when biennial and perennial species are in the seedling to rosette stage and before flower stalks appear. For difficult to control perennial broadleaf weeds and woody species, use up to 1 gallon of this product and 1 to 4 quarts of a herbicide containing triclopyr, triethylamine salt per acre.
Biennial and perennial broadleaf weeds and susceptible woody plants	2.75 to 5.5 pints	For conifer release, make application in early spring before budbreak of conifers when weeds are small and actively growing.
Spot treatment to control broadleaf weeds	1.0 fl oz/ gal of spray solution (see instructions for Spot Treatment)	To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the specified broadcast rate and spray to thoroughly wet all foliage. Mix 1.0 fluid ounces per gallon of spray solution and apply through pump up sprayer or backpack sprayer. Addition of a non-ionic surfactant is recommended to improve coverage. See rate conversion table and instructions for Spot Treatment and use of hand-held sprayers under Application Directions.
Conifer release: species: Balsam fir, Black spruce, Jack pine, Ponderosa pine, Red pine, Red spruce, White pine, White spruce	2 to 4 pints	To control competing hardwood species, alder, aspen, birch, hazel, and willow, apply from mid to late summer when growth of conifer trees has hardened off and woody plants are still actively growing. Apply with ground or air equipment, using sufficient spray volume to ensure complete coverage. Because this treatment may cause occasional conifer injury, do not apply if such injury cannot be tolerated.
Directed spray: Conifer plantations including pine	5.5 pints per 100 gallons	Apply when brush or weeds are actively growing by directing the spray so as to avoid contact with conifer foliage and injurious amounts of spray. Apply in oil, oil-water, or water carrier in a spray volume of 10 to 100 gallons per acre, or sufficient spray solution for adequate coverage.
Basal spray	5.5 quarts per 100 gallons or 1.75 fl oz/gal of water	Thoroughly wet the base and root collar of all stems until the spray begins to accumulate around the root collar at the ground line. Wetting stems with the mixture may also aid in control.
Surface of cut stumps		Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and girdle		Cut frills (overlapping, V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Treat freshly cut frills with as much of the 2,4-D mixture as they will hold.

Tree injection	0.025 to 0.05 fl oz (0.75 to 1.5 mL) per injection site	To control unwanted hardwood trees, elm, hickory, oak, and sweetgum, in forests and other non-crop areas, apply by injecting at a rate of 0.034 fl oz (1 mL) of undiluted BCB 2,4-D LV6 per inch of trunk diameter at breast height (DBH) as measured approximately 4 1/2 foot above the ground. However, injection should occur as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Do not treat maples during the spring sap flow. For hard to control species, ash, maple, and dogwood, use 0.05 fl oz (1.5 mL) of undiluted BCB 2,4-D LV6 per injection site or double the number of 3/4 ml injections. No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.
-----------------------	---	---

RESTRICTIONS FOR USE ON FORESTRY

- Do not allow sprays to contact conifer shoot growth (current year's new growth) or injury may occur.
- Do not apply to nursery seed beds.
- For conifer release, do not use on plantations where pine or larch are among the desired species.
- For broadcast applications, do not apply more than a total of 5.8 pints (4.0 lb 2,4-D ae) per acre per 12-month period.
- Limited to one broadcast application, one basal spray or cut surface application, or one injection application per year.
- For aerial application, use a minimum of 2 gallons of water per acre.
- For basal spray, cut surface stumps, and frill applications, do not apply more than 11.6 pints (8.0 lbs of 2,4-D ae) per 100 gallons of spray solution.
- For tree injections: Do not apply more than 2 ml of this product per injection site. Limited to 1 injection application per year.

FOREST MANAGEMENT

CONIFER RELEASE

To control Alder, apply 2 to 2.75 pints of this product in 8 to 25 gallons water per acre as a foliage spray, or sufficient spray solution for adequate coverage. Treat when 3/4 of the brush foliage has attained full size leaves and before new conifer growth reaches 2 inches in length. This is usually between early May and mid-June. Adjust treatment date depending on stage of growth of conifers and brush species. This may cause leader deformation on exposed firs, but they should overcome this during the second year after spraying.

To control susceptible trees and brush species, Ceanothus spp., Chinquapin, Madrone, Manzanita, Oak and Tanoak and to release Douglas fir, Grand fir, Hemlock, or Sitka spruce, apply 2 quarts of product per acre before new growth on Douglas fir is 2 inches long. To control Manzanita and Ceanothus in Ponderosa pine, apply 2 quarts of this product before pine growth begins in spring.

To increase performance, add 2 to 4 quarts of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at label rate.

After Black spruce, Jack pine, Northern conifers, Red pine and White spruce cease growth and "harden off" (usually in mid-July), a spray of 1 to 2 quarts of product in 8 to 25 gallons of water per acre, or sufficient spray solution for adequate coverage. Product may be applied by air to control certain competing hardwood species, Alder, Aspen, Birch, and Willow. Since this treatment may cause occasional conifer injury, do not use if such injury cannot be tolerated. Consult your Regional or Extension Forester or State herbicide specialist for recommendations to fit local conditions.

To control Hazel brush in the Lake states, apply 2.75 pints of this product in 6 to 25 gallons of water per acre, or sufficient spray solution for adequate coverage. Apply when new shoot growth of Hazel is complete (usually mid-July).

Tree Injections (Pine Release): To control hardwoods, Elm, Hawthorn, Hickory, Maple, Oaks, Pecan, Sumac and Sweetgum in forest and other non-crop areas, apply this product undiluted in a concentrate tree injector calibrated to apply 1.0 ml per injection. Space injections 2 inches apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species, Ash, Blue beech, Dogwood, Hickory, and Red maple, make injections 1 to 1.5 inches apart, edge to edge. Treatment may be made at any time of the year. For best results, injections should be made during growing season, May 15 to October 15. For dilute injections, mix 5-1/2 pints of this product in 19 gallons of water. No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

Dormant Application (other than pine): For the control of susceptible deciduous brush species, Alder, Cascara, Cherry poplar and Serviceberry, apply up to 2 quarts of product per acre in sufficient diesel, fuel oil or kerosene for good coverage.

Application may be made by ground or air and should be made before conifer bud break.

Pine Only: Make application while pine buds are still dormant. Apply 2.75 pints of product per acre in sufficient spray solution for adequate coverage by air or ground equipment. Do not use this application unless some pine injury is acceptable. Use of diesel, kerosene, or other oil, or addition of surfactants to spray mix may cause unacceptable pine injury.

Christmas Tree Plantations: For control of labeled broadleaf weeds in Douglas Fir Christmas trees, use 0.75 to 1.33 pints of this product per acre.

Apply over the top of Douglas Fir by ground or aerial application equipment only when the trees are dormant, prior to bud break. Do not spray over the top of pine or true firs (*Abies* spp.).

Directed Sprays may be made to weeds in Christmas tree plantations of all conifer species, but the spray must not contact tree foliage as injury may occur. Do not apply to weakened, diseased, or stressed seedlings since unacceptable injury can occur. This product may be mixed with Atrazine for Christmas tree application. (See Tank Mix section.)

Herbaceous Weed Control: To control over-wintering susceptible weeds, False dandelion, Klamath weed, Plantain, Tansy ragwort, apply 1.33 to 4 pints of product in sufficient spray solution for adequate coverage. Make application at rates and timing indicated above if Pines are present. For control of Hazel brush and similar species in the Lake States area, apply 2.75 pints of product per acre in 8 to 25 gallons of water when new shoot growth of Hazel is complete (usually mid-July).

SITE PREPARATION

Budbreak Spray: To control Alder and other susceptible species before planting forest seedlings, apply 2.75 to 5.5 pints of this product in 8 to 25 gallons fuel oil per acre, or sufficient spray solution for adequate coverage. Apply after alder buds break, but before foliage is 1/4 full size. Application may be made by air or ground. If desired, water, diesel, or kerosene may be substituted for fuel oil as diluent.

Foliage Spray: To control Alder before planting forest tree seedlings, apply 2.75 pints of this product plus 2 quarts fuel oil in 8 to 25 gallons of water per acre, or sufficient spray solution for adequate coverage. Apply after most Alder leaves are full size. To increase penetration, 2 to 4 quarts per acre of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at label rates may be added to the spray mixture.

The maximum rate per broadcast application is 5.8 pints (4 lbs 2,4-D ae) per acre, limited to one application per year.

Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

RANGELANDS AND ESTABLISHED GRASS PASTURES

Included Perennial Grasslands not in Agricultural Production, Such as Conservation Reserve Program Acres

TARGET WEEDS OR WOODY PLANT	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Annual broadleaf weeds Perennial and biennial broadleaf weeds	1.33 pints 1.33 to 2.75 pints	For best results, apply when weeds are small and actively growing, before the bud stage. Apply when musk thistles or other biennial species are in the seedling to rosette stage and before flower stalks appear. Refer to the Weeds Controlled section for a listing of susceptible weed species and weeds that may be only partially controlled and require repeat applications and/or use of higher specified rates, even under ideal conditions of application
Spot treatment to control broadleaf weeds	1.0 fl oz/ gal of spray solution (see instructions for Spot Treatment)	To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the specified broadcast rate and spray to thoroughly wet all foliage. Mix 1.0 fluid ounces per gallon of spray solution and apply through pump up sprayer or backpack sprayer. The addition of a non-ionic surfactant is recommended to improve coverage. See rate conversion table and instructions for Spot Treatment and use of hand-held sprayers under Application Directions.
Tree injection	0.75 to 1.5 mL per injection site	To control unwanted hardwood trees, elm, hickory, oak, and sweetgum, in forests and other non-crop areas, apply by injecting at a rate of 1 mL of undiluted BCB 2,4-D LV6 per inch of trunk diameter at breast height (DBH) as measured approximately 4 1/2 foot above the ground. However, injection should occur as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Do not treat maples during the spring sap flow. For hard to control species, ash, maple, and dogwood, use 1.5 mL of undiluted BCB 2,4-D LV6 per injection site [or double the number of 0.75 ml injections]. No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.
Wild garlic and wild onion	2.75 pints	Make three applications (fall-spring-fall or spring-fall-spring) starting in late fall or early spring.
Broadleaf weed control in newly sprigged coastal bermudagrass	1.33 to 2.75 pints	Applications may be made either preemergence or postemergence. Follow Specific Use Directions for annual, biennial and perennial broadleaf weed control, above.
Sand shinnery oak, and Sand sagebrush	1.33 pints	Sand shinnery oak: Apply by aircraft between May 15 and June 15. Sand sagebrush: Apply by ground or aircraft when foliage is fully expanded and plants are actively growing.
Big sagebrush, rabbitbrush, chamise, manzanita, buckbrush, coastal sage, coyotebrush, and chaparral species.	2.75 pints	Apply by ground or aircraft when foliage is fully expanded and plants are actively growing. Re-treatment may be needed.

TARGET WEEDS OR WOODY PLANT	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Southern wild rose Broadcast application Spot treatment	2.75 pints 1.0 fl oz/ gal of spray solution	Broadcast: Apply in a spray volume of 2 gallons or more per acre by aircraft or 10 gallons or more per acre by ground equipment, or sufficient spray solution for adequate coverage. Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Mix 1.0 fl oz per gallon of spray solution and apply through pump up sprayer or backpack sprayer. Addition of a non-ionic surfactant is recommended to improve coverage. Two or more treatments may be required. Do not exceed 2.75 pints per acre per application.
Surface of cut stumps	1.75 fl oz/gal of water	Apply as soon as possible after cutting trees. Thoroughly soak the entire stump with the 2,4-D mixture including cut surface, bark and exposed roots.
Frill and girdle		Cut frills (overlapping, V-shaped notches cut downward through the bark in a continuous ring around the base of the tree) using an axe or other suitable tool. Treat freshly cut frills with as much of the 2,4-D mixture as they will hold.

PRECAUTIONS FOR USE IN RANGELAND AND PASTURE

- For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

RESTRICTIONS FOR USE ON RANGELAND AND PASTURE

- The preharvest interval (PHI) to forage hay is 7 days. For program lands, such as CRP, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.
- If grass is to be cut for hay, Agricultural Use Requirements for the Worker Protection Standard are applicable.
- Do not use on bentgrass, alfalfa, clover, or other legumes.
- Do not use on newly seeded areas until grass is well established.
- Do not use from early boot to milk stage where grass seed production is desired.
- Do not apply within 30 days of a previous application.
- For grazed areas, the maximum use rate is 2.9 pints of **BCB 2,4-D LV6** (2.0 lbs of 2,4-D ae) per acre per application.
- Do not apply more than a total of 5.8 pints of **BCB 2,4-D LV6** (4.0 lbs of 2,4-D ae) per acre per year.
- Do not make more than two applications per year.
- For susceptible annual and biennial broadleaf weeds: Do not apply more than 1.5 pints (1.0 lb of 2,4-D ae) per acre per application.
- For moderately susceptible biennial, perennial broadleaf weeds and difficult to control weeds and woody plants: Do not apply more than 2.9 pints (2.0 lbs of 2,4-D ae) per acre per application.
- **Spot treatment:** Do not apply more than 2.9 pints (2.0 lbs of 2,4-D ae) per acre.

NON-CROPLAND AREAS

Fencerows, Hedgerows, Roadsides, Ditches, Rights-of-Way, Utility Power Lines, Railroad, Airports, Airfields, Vacant Lots, Highway and Industrial Sites

TREATMENT SITE / METHOD OF APPLICATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Annual broadleaf weeds	1.33 to 2.75 pints	Apply when annual weeds are small and growing actively before the bud stage. Biennial and perennial weeds should be rosette to bud stage, but not flowering at the time of application. For difficult to control perennial broadleaf weeds and woody species, tank mix up to 5.5 pints of BCB 2,4-D LV6 plus an appropriately labeled herbicide containing triclopyr, triethylamine salt at the specified rate per acre. For ground application: High volume - apply a total of 100 to 400 gallons per acre; low volume - apply a total of 10 to 100 gallons per acre, or sufficient spray solution for adequate coverage. For helicopter: Apply a total of 2 to 30 gallons per acre spray volume, or sufficient spray solution for adequate coverage.
Perennial and Biennial broadleaf weeds	2.75 pints	
Susceptible woody plants on rights-of-way	2.75 to 5.5 pints	
Spot treatment to control broadleaf weeds	1.0 fl oz/ gal of spray solution (see instructions for Spot Treatment)	To control broadleaf weeds in small areas with a hand sprayer, use an application rate equivalent to the specified broadcast rate and spray to thoroughly wet all foliage. Mix 1.0 fluid ounces per gallon of spray solution and apply through pump up sprayer or backpack sprayer. Addition of a non-ionic surfactant is recommended to improve coverage. See rate conversion table and instructions for Spot Treatment and use of hand-held sprayers under Application Directions.
Tree injection	0.75 to 1.5 mL per injection site	To control unwanted hardwood trees, elm, hickory, oak, and sweetgum, in forests and other non-crop areas, apply by injecting at a rate of 1 mL of undiluted BCB 2,4-D LV6 per inch of trunk diameter at breast height (DBH) as measured approximately 4 1/2 foot above the ground. However, injection should occur as close to the root collar as possible and the injection bit must penetrate the inner bark. Applications may be made throughout the year, but for best results apply between May 15 and October 15. Do not treat maples during the spring sap flow. For hard to control species, ash, maple, and dogwood, use 1.5 mL of undiluted BCB 2,4-D LV6 per injection site or double the number of 0.75 ml injections. No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.
Southern wild rose Broadcast application	2.75 pints plus 4 to 8 oz.	Broadcast: Use up to 5.5 pints of this product plus 4 to 8 ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well

TREATMENT SITE / METHOD OF APPLICATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Spot treatment	agricultural surfactant 1.0 fl oz/ gal of spray solution	developed. Two or more treatments may be required Spot treatment: Apply when foliage is well developed. Thorough coverage is required. Mix 1.0 fl oz per gallon of spray solution and apply through pump up sprayer or backpack sprayer. The addition of a non-ionic surfactant is recommended to improve coverage. Two or more treatments may be required.

PRECAUTIONS FOR USE IN NON-CROPLAND

- Bentgrass, St. Augustine, clover, legumes and dichondra may be severely injured or killed by this treatment.
- Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial uses, or for commercial seed production, or for research purposes.

RESTRICTIONS FOR USE ON NON-CROPLAND

- Do not apply to newly seeded areas until grass is well established.
- Annual and perennial weeds: Do not apply more than 2.9 pints of **BCB 2,4-D LV6** (2.0 lbs of 2,4-D ae) per acre per application.
Do not make more than two applications per year.
Do not reapply to a treated area within 30 days of a previous application.
- Woody plants:
Do not apply more than a total of 5.8 pints of pints of **BCB 2,4-D LV6** (4.0 lbs of 2,4-D ae) per acre per year.
Do not make more than one application per year.

TANK MIXTURES IN NON-CROP AREAS

**BCB 2,4-D LV6 and Products Containing Triclopyr, butoxyethyl ester or Triclopyr, triethylamine salt
Tank Mixtures* for Non-Crop Areas**

WEEDS IN CROP	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Broadleaf weed control	1.33 to 2.75 pints BCB 2,4-D LV6 plus Triclopyr, butoxyethyl ester or Triclopyr, triethylamine salt at specified label rates	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 2 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing.
Woody plant control Broadcast foliar spray	5.5 pints BCB 2,4-D LV6 plus Triclopyr, butoxyethyl ester or Triclopyr, triethylamine	Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons of water per acre. Apply when woody plants are actively growing.

	salt at specified label rates	
Woody plant control High volume leaf-stem treatment with ground equipment	1.33 to 5.5 pints BCB 2,4-D LV6 plus Triclopyr, butoxyethyl ester or Triclopyr, triethylamine salt at specified label rates	Mix 1.33 to 2.75 pints BCB 2,4-D LV6 plus 1.5 to 3 pints (0.75 to 1.5 lbs) triclopyr, butoxyethyl ester or 2 to 4 pints (0.75 to 1.5 lbs) triclopyr, triethylamine salt in a mixture of water volume intended per acre. This may require up to 100 to 400 gallons of water per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems and root collars of plants to be controlled.
Woody plant control Aerial application (Helicopter Only)	5.5 pints BCB 2,4-D LV6 plus Triclopyr, butoxyethyl ester or Triclopyr, triethylamine salt at specified label rates	Apply in a total mixture of water volume intended per acre. This may require a volume of 10 to 30 gallons per acre using drift control equipment such as the MICRO-FOIL® boom or an effective drift control agent. Use the higher rates and volumes when plants are dense or under drought conditions.

*It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

TANK MIXTURES IN NON-CROP AREAS

BCB 2,4-D LV6 and Products Containing Dicamba, diethylamine salt Tank Mixtures* for Non-Crop Areas

WEEDS IN CROP	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Annual broadleaf control	1.33 to 2.75 pints BCB 2,4-D LV6 plus dicamba, diethylamine salt at specified label rates	For wider spectrum control of broadleaf weeds and woody plants: Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre, or sufficient spray solution for adequate coverage. Apply when broadleaf weeds are actively growing.
Perennial and biennial broadleaf weeds	2 to 2.75 pints BCB 2,4-D LV6 plus dicamba, diethylamine salt at specified label rates	Apply as a broadcast spray which may require 20 to 100 gallons total spray per acre, or sufficient spray solution for adequate coverage. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3-inch rosette stage.

Woody plant control Broadcast, high volume, stem foliage or aerial application	5.5 pints BCB 2,4-D LV6 plus dicamba, diethylamine salt at specified label rates	Apply as a broadcast spray in a mixture of water volume intended per acre or as a high volume stem foliage spray in enough volume (20 to 100 gallons of water) to thoroughly wet leaves, stems and root collars or apply aerially in enough water to deliver total spray volume of 10 to 30 gallons of water per acre using drift control agent, or sufficient spray solution for adequate coverage. This may require 100 to 400 gallons of water per acre for adequate coverage. Use the higher rates and volumes when plants are dense or under drought conditions.
--	---	---

*It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

SMALL AREA APPLICATIONS FOR NON-CROP USE

For control of broadleaf weeds in small non-crop areas with hand held or back-pack sprayers mix 2-3/4 fluid ounces of this product per gallon of water. Thoroughly wet all weed foliage. Maintain agitation of mixture to prevent separation.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

TURFGRASS GROWN FOR SEED OR SOD FARMS

TREATMENT SITE / METHOD OF APPLICATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Grasses grown for seed (postemergence use) Seedling grass (five-leaf stage or later) Well-established grasses	0.5 to 0.75 pint 0.75 to 2.75 pints	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth. Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 0.75 pint per acre. Cool season grasses are tolerant of higher rates. Do not apply to grass in the early boot through milk stage if seed production is desired. When grass is well established, higher rates of up to 2.75 pints per acre may be applied for control of hard to kill annual or perennial weeds.
Sod farms (postemergence)	1.33 to 2.75 pints	Deep-rooted perennials, bindweed and Canada thistle may require repeat applications. Avoid mowing sod farms for 1 to 2 days before or after application. Delay irrigation until the day following application.

PRECAUTIONS FOR USE IN TURFGRASS GROWN FOR SEED OR SOD

- **Reseeding:** Delay reseeding at least 30 days following application. Preferably, with spring application, reseed in the fall and with fall application, reseed in the spring.

RESTRICTIONS FOR USE IN TURFGRASS GROWN FOR SEED OR SOD

- The preharvest interval (PHI) is 7 days.
- Do not apply more than a total of 5.8 pints of **BCB 2,4-D LV6** (4.0 lbs of 2,4-D ae) per acre per year, limited to two applications per year.

- The maximum rate per broadcast application is 2.9 pints **BCB 2,4-D LV6** (2.0 lbs of 2,4-D ae) per acre.
- Do not use on creeping grasses such as bent except as a spot treatment.
- Do not use on injury-sensitive southern grasses, such as St. Augustinegrass.
- Do not use on dichondra or other herbaceous groundcovers. Legumes may be damaged or killed.
- Minimum of 21 days between applications.

**Ornamental Turfgrass (Excluding Grasses Grown for Seed or Sod Farms)
Includes Lawns, Golf Courses (Aprons, Fairways, Roughs and Tees), Cemeteries,
Parks, Sports Fields**

TREATMENT SITE / METHOD OF APPLICATION	AMOUNT OF BCB 2,4-D LV6 PER ACRE	DIRECTIONS
Ornamental turfgrass (postemergence use)		
Seedling grass (five-leaf stage or later)	0.5 to 0.75 pint	Apply when weeds are small and actively growing. For best results, apply when soil moisture is adequate for active weed growth.
Well-established grasses	1.33 to 2 pints	Deep-rooted perennial weeds, bindweed and Canada thistle may require repeat applications.
Biennial and perennial broadleaf weeds	2 pints	Do not apply to newly seeded grasses until well established (five-leaf stage or later) and then use a maximum of 0.75 pint per acre. Cool season grasses are tolerant of higher rates.

PRECAUTIONS FOR USE IN ORNAMENTAL TURFGRASS

- **Reseeding:** Delay reseeding at least 30 days following application. Preferably, with spring application, reseed in the fall and with fall application, reseed in the spring.

RESTRICTIONS FOR USE IN ORNAMENTAL TURFGRASS

- Do not make more than two broadcast applications per year per treatment site (does not include spot treatments).
- Do not apply more than a total of 4.3 pints of **BCB 2,4-D LV6** (3.0 lbs of 2,4-D ae) per acre per year.
- Do not use on creeping grasses such as bent except as a spot treatment.
- Do not use on injury-sensitive southern grasses, such as St. Augustinegrass.
- Do not use on dichondra or other herbaceous groundcovers. Legumes may be damaged or killed.
- Minimum of 21 days between applications.

TANK MIXES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions, limitations, and directions for use on all product labels involved in the tank mixture. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Using this product and products containing bromoxynil for weed control on cereal grains (barley, rye, wheat, and oats): Using products containing bromoxynil octanoate (BO) at specified label rates will control some annual weeds that are resistant to this product and may be tank mixed with this product for broader spectrum weed control on small grains. In cereal areas except Idaho, Oregon, Colorado, Wyoming, Montana, and Washington, use 0.33 to 0.75 pint of this product plus an appropriately labeled bromoxynil product at specified label rates. In Idaho, Oregon, Colorado, Wyoming, Montana, and Washington, use 0.33 to 0.75 pint of this product plus an appropriately labeled bromoxynil product at specified label rates. First mix this product in water, then add the bromoxynil octanoate. Use the higher rates for larger weeds or where weed growth is slow due to dry or cold weather. Apply before weeds are 6 inches high. Use 10 to 25 gallons total spray volume per acre with ground equipment or 2 to 10 gallons total spray volume with air application, or sufficient spray solution for adequate coverage. Use higher volume on larger weeds.

Using this product with products containing dicamba salts to provide more complete Kochia

control: Offers quick burndown. Provides residual activity to control later weed flushes making harvesting easier and reducing postharvest weed control needs. Controls broader weed spectrum while offering better control of Flixweed, Mustards, Russian thistle, and Wild buckwheat. Controls large weeds. Allows for early treatment. Apply 5.5 ounces (0.25 lb 2,4-D a.e.) of this product with a product containing metsulfuron plus a product containing dicamba diethylamine or sodium salt as the active ingredient at the specified label rates per area. The tank mix can be applied to Winter wheat and the four-leaf stage (tillering) to prior to joint. It can be applied to Spring wheat from the four-leaf stage through the five-leaf stage. Growers who want to rotate to a sensitive crop following wheat and are concerned about carryover from a metsulfuron application can substitute a product containing tribenuron-methyl in the tank mix which allows crop rotation 60 days after application. Use products containing tribenuron-methyl at the specified labeling use rate per acre.

Using this product with products containing metrizubin as knockdown herbicides for no till: This product with products containing the active ingredient metrizubin may be applied as an early preplant surface application for the control of certain broadleaf weeds and grasses in soybeans in minimum or no-till products. Application is recommended 30 days prior to planting. Apply at rate of 1.33 pints of this product (1 lb 2,4-D a.e.) per acre with labeled rates specified by the metrizubin product label. Where grass herbicide is used in tank mix, apply at the rates specified on that product's label.

Using this product with products containing atrazine for weed control in Christmas tree and forest plantings: A tank mix of these two products can be used to control weeds and thus aid in the establishment of young transplants of Austrian pine, Bishop pine, Blue spruce, Douglas fir, Grand fir, Jeffrey pine, Knobcone pine, Loblolly pine, Lodgepole pine, Monterey pine, Nobel fir, Ponderosa pine, Scotch pine, Sitka spruce, Slash pine, and White fir.

The mix should be applied between fall and early spring, preferably in February or March, while trees are still dormant, or soon after transplanting. Weeds should not be more than 1.5 inches high. It can be applied with either ground or air equipment. Helicopters have been highly effective for reforestation applications or steep terrain. Uniform application is the key to good weed control. Use 10 to 40 gallons of water per acre for ground applications; a minimum of 2 gallons of water when applying by air, or sufficient spray solution for adequate coverage. Be sure equipment is properly calibrated. All screens in the spray system -- nozzles, and in-line and suction strainers -- should be 15 mesh or coarser. Use a pump with capacity to maintain a nozzle pressure of 35 to 40 psi, and sufficient agitation to keep the mixture in suspension in the spray tank. If a nurse tank is used, keep the mixture agitated while awaiting transfer to the spray tank. Mix a product containing atrazine at the specified label rate with 1.5 to 4 pints (1 to 3 lbs 2,4-D ae) of this product. The actual rate of the product containing atrazine must depend on soil type. Soils high in organic matter require higher rates than light to medium soils. Band application to Christmas Trees - Calculate the amount to be applied per acre. The band width in inches, divided by the rows spacing in inches, times the rate per acre for broadcast treatment will equal the amount needed per acre for band treatment. Please read the product label(s) for additional instructions.

Using this product with products containing flumioxazin in reduced-tillage or no-till systems: This product may be applied in combination with a product containing the active ingredient flumioxazin for the control of annual grasses and broadleaf weeds and the suppression of emerged perennial weeds when soybeans are directly seeded into a stale seedbed, cover crop or in previous crop residues. Special precautions: Poor weed control and/or crop injury may result if directions are not followed. Do not use a rib-type press wheel on your no-till planter or crop injury may result. Apply at a rate of 1.33 pints of this product (1 lb 2,4-D ae) per acre with labeled rates of herbicide containing flumioxazin. Application is recommended 30 days prior to planting.

Using this product with products containing sethoxydim as a burndown prior to planting soybeans: For broad spectrum post-emergence weed control, a tank mix application of this product with a product containing the active ingredient sethoxydim may be made for control of emerged broadleaf and grass weeds before planting soybeans. Apply at a rate of 0.75 pint of this product (0.5 lb 2,4-D ae) per acre with labeled rates of the product containing sethoxydim.

Using this product with products containing imazaquin in preplant applications in no-till soybeans: For broad spectrum post-emergence weed control, a tank mix application of this product with products

containing the active ingredient imazaquin may be made for the control of emerged broadleaf and grass weeds before planting soybeans. Apply at a rate of 0.75 pint of this product (0.5 lb 2,4-D ae) per acre up to 7 days prior to planting, or 1.33 pints (1 lb 2,4-D ae) per acre up to 30 days prior to planting, with labeled rates of the herbicide containing imazaquin.

Using this product with products containing triclopyr, butoxyethyl ester or triclopyr, triethylamine salt for non-crop areas: See **TANK MIXTURES IN NON-CROP AREAS BCB 2,4-D LV6 and Products Containing Triclopyr, butoxyethyl ester or Triclopyr, triethylamine salt Tank Mixtures for Non-Crop Areas** section of this label.

Using this product with products containing dicamba, diethylamine salt tank mixtures for non-crop areas: See **TANK MIXTURES IN NON-CROP AREAS BCB 2,4-D LV6 and Products Containing Dicamba, diethylamine salt Tank Mixtures for Non-Crop Areas** section of this label.

Using this product with products containing metsulfuron methyl or sulfometuron methyl: To improve control of some target species, this product may also be tank mixed with herbicides containing metsulfuron methyl or sulfometuron methyl for postemergent weed control. Tank mixes have shown improved control where resistant bio-types are present.

NOTE: All intended tank mix combinations should be used only in areas specified on the same broadleaf weed species found on both labels.

For application methods and other use specifications, use the most restricted limitations from labeling of both products.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a dry, secured storage area. Keep container tightly closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING:

[Nonrefillable Containers 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

[Nonrefillable Containers Larger than 5 Gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30

seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times.

Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

[Refillable Containers Larger than 5 Gallons:] Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of BETTER CHOICE BRANDS LLC or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold BETTER CHOICE BRANDS LLC and Seller harmless for any claims relating to such factors.

BETTER CHOICE BRANDS LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to or beyond the control of Seller or BETTER CHOICE BRANDS LLC and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, BETTER CHOICE BRANDS LLC MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, neither BETTER CHOICE BRANDS LLC nor Seller shall be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF BETTER CHOICE BRANDS LLC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF BETTER CHOICE BRANDS LLC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

BETTER CHOICE BRANDS LLC and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of BETTER CHOICE BRANDS LLC.

